

Applicant : Katsuna [REDACTED] et al.
Serial No. : 09/830,548
Filed : April 26, 2001
Page : 3

Attorney's Doc : 06501-077001 / J1-A0004P-US

REMARKS

Applicants hereby submit that the enclosures fulfill the requirements under 37 C.F.R. §1.821-1.825. The amendments in the specification merely insert the paper copy of the Sequence Listing and sequence identifiers in the specification. No new matter has been added.


Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment.

Please apply any charges or credits to Deposit Account No. 06-1050.

Respectfully submitted,

Date:

Aug. 29, 2001



Janis K. Fraser, Ph.D., J.D.
Reg. No. 34,819

Fish & Richardson P.C.
225 Franklin Street
Boston, Massachusetts 02110-2804
Telephone: (617) 542-5070
Facsimile: (617) 542-8906

“Version With Markings to Show Changes Made”

In the specification:

Paragraph beginning at page 4, line 18. has been amended as follows:

The ligands for CD28 and CTLA-4 are CD80 (B7-1) and CD86 (B7-2) in human and mice. CTLA-4 has about 20 times as high affinity to both ligands as CD28. It has been elucidated that the amino acid sequence structures “MYPPPY (Met-Tyr-Pro-Pro-Pro-Tyr; SEQ ID NO:1)” conserved through animal species is important for the binding of CD28 and CTLA-4 to CD80 (B7-1). It has also been reported that, when CD28 is stimulated, PI3 kinase (phosphoinositide 3 kinase, PI3K) associates with the phosphorylated tyrosine residue in a partial sequence “YMNM (Tyr-Met-Asn-Met; SEQ ID NO:2)” of CD28 and that CD28 plays an important role in intracellular signal transmission through this “YxxM” structure. Furthermore, it has been reported that CTLA-4 also has a sequence represented by “YxxM,” namely “YVKM (Tyr-Val-Lys-Met; SEQ ID NO:3)” in its cytoplasmic region and that, after being stimulated, SYP associates with this sequence.